arXiv:1401.5601v1 [math.CO] 22 Jan 2014

# Unimodality and genus distributions

**Liangxia Wan** *

*Department of Mathematics, Beijing Jiaotong University, Beijing* 100044*, China*

**Abstract**– New criteria are shown that certain combinations of finite unimodal polynomials are unimodal. As applications, unimodality of several polynomial sequences satisfying dependent recurrence relations and their modes are provided. Then unimodality of genus distributions for some ladders and crosses can be determined. As special cases, that of genus distributions for Closed-end ladders, circular ladders, Möbius ladders and Ringel ladders and their modes are given, which induces the known results for Closed-end ladders.

**Keywords**– Unimodal, log-concave, genus distribution, embedding

## 1. Introduction

Let $S_i$, $\gamma$ and $\gamma_M$ denote the surface of genus $i$, the minimum genus and maximum genus of a graph $G$ respectively. Let $g_i(G)$ denote the number of 2-cell embeddings of $G$ embedded on $S_i$ for $i \geq 0$. The *genus distribution* of $G$ is a sequence of numbers followed:

$$g_\gamma(G), g_{\gamma+1}(G), g_{\gamma+2}(G), \cdots, g_{\gamma_M}(G).$$

$f_G(x) = \sum_{i=\gamma}^{\gamma_M} g_i(G)x^i$ is called the *genus polynomial* of $G$. In general, it is an NP-complete problem to determine the genus distribution of a graph, since evaluating $\gamma$ is an NP-complete problem [12].

A finite sequence of nonnegative numbers $\{a_i\}_{i=q}^n$ is *unimodal* if there exist index $q \leq l \leq m \leq n$ such that $a_q \leq a_{q+1} \leq \cdots \leq a_{l-1} < a_l = \cdots = a_m > a_{m+1} \geq \cdots \geq a_n$ where integers $0 \leq q \leq n$. The corresponding polynomial $\sum_{i=1}^{n} a_i x^i$ is also called *unimodal*. $l, l+1, \cdots, m$ are called the *modes* of the sequence. If $l = m$, then $m$ is called a *peak* of the sequence. The sequence is *log-concave* if $a_i^2 \geq a_{i-1}a_{i+1}$ for $1 \leq i \leq n-1$. Obviously, if a sequence is log-concave, then it is unimodal. Unimodality problems arise naturally in many branches of mathematics and have been extensively investigated. See Stanleys [10] and Brentis survey articles [1] for details. Throughout this paper each sequence of numbers is that of nonnegative numbers. $[a]$ denote the maximum integer not more than $a$ for a nonnegative number $a$.

*E-mail address: lxwan@bjtu.edu.cn. *Partially supported by NNSFC under Grants No.* 11201024

Unimodality problems for the genus distribution of a graph include unimodality, log-concavity and the reality of zeros of a genus polynomial. Gross, Robbins and Tucker proposed the conjecture that the genus distribution of every graph is log-concave and showed that genus distributions of bouquets are log-concave [5]. Furst, Gross and Statman showed that genus distributions of Closed-end ladders and cobblestone paths are log-concave [3]. Gross, Mansour and Tucker proved genus distributions of some ring-like graphs are log-concave [4]. Stahl proved that all zeros of genus polynomials of tree-like graphs, cobblestone paths, diamond bands and some vertex-forest multijoins are real and negative [9]. Chen, Liu and Wang also considered zeros of genus polynomials of some graphs [7, 2]. In 2003, Liu conjectured that the genus distribution of each graph is unimodal [6]. Zhao and Liu verified that the genus distribution of every tree graph is unimodal [17].

This paper mainly concerns unimodality and log-concavity of polynomial sequences there exist dependent recurrence relations among them. In Section 2, new criteria are presented to determine unimodality of a sequence of numbers generated from finite unimodal sequences of numbers. In Section 3, genus distributions for sets of ladder surfaces are shown to be unimodal or even log-concave. Then unimodality of genus distributions for some ladders and crosses can be verified. As special consequences unimodality of genus distributions for Closed-end ladders, circular ladders, Möbius ladders [8] and Ringel ladders [11] are proved and their modes of genus distributions for these graphs are determined in Section 4, which induces the known results for Closed-end ladders. Section 5 gives some problems for further study.

## 2. Criteria

**Theorem** 2.1 *Suppose that* $\{x_i\}_{i=q_1}^{n_1}$ *and* $\{y_i\}_{i=q_2}^{n_2}$ *are unimodal sequences of numbers for* $0 \leq q_1 \leq n_1$ *and* $0 \leq q_2 \leq n_2$. *Let* $l_1, l_1+1, \cdots, m_1$ *and* $l_2, l_2+1, \cdots, m_2$ *be the modes of* $\{x_i\}_{i=q_1}^{n_1}$ *and* $\{y_i\}_{i=q_2}^{n_2}$ *for* $j = 1, 2$ *and* $l_j \leq m_j$ *respectively, let* $r_j$ *be non-negative integers and let* $a_j > 0$. *Then the sequence* $\{a_1x_{i-r_1} + a_2y_{i-r_2}\}$ *is unimodal if and only if it is unimodal for* $\min\{l_1+r_1, l_2+r_2\} \leq i \leq \max\{m_1+r_1, m_2+r_2\}$.

**Proof.** Put $z_i = a_1x_{i-r_1}+a_2y_{i-r_2}$. Clearly, the sequence $\{z_i\} = \{z_i\}_{i=q}^{n}$ where $q = \min\limits_{1\leq j\leq 2}\{q_j+r_j\}$ and $n = \max\limits_{1\leq j\leq 2}\{n_j+r_j\}$. For brevity, it is denoted by $\{z_i\}$. Put $l = \min\{l_1+r_1, l_2+r_2\}$ and put $m = \max\{m_1+r_1, m_2+r_2\}$. Since $\{x_i\}_{i=p_1}^{n_1}$ and $\{y_i\}_{i=p_2}^{n_2}$ are unimodal and since $a_j > 0$ for $j = 1, 2$, these follow that

$$z_q \leq z_{q+1} \leq \cdots \leq z_{l-1} < z_l = z_m > z_{m+1} \geq \cdots \geq z_n.$$

Thus the result is obvious. □

If $m - l \leq 3$, then it is obvious that $\{a_1x_{i-r_1} + a_2x_{i-r_2}\}$ is unimodal for $l \leq i \leq m$ and so the following conclusion holds.

**Corollary** 2.2 *Suppose that* $\{x_i\}_{i=q_1}^{n_1}$ *and* $\{y_i\}_{i=q_2}^{n_2}$ *are unimodal sequences of numbers for* $0 \leq q_1 \leq n_1$ *and* $0 \leq q_2 \leq n_2$. *Let* $l_1, l_1+1, \cdots, m_1$ *and* $l_2, l_2+1, \cdots, m_2$ *be the modes of* $\{x_i\}_{i=q_1}^{n_1}$ *and* $\{y_i\}_{i=q_2}^{n_2}$ *for* $j = 1, 2$ *and* $l_j \leq m_j$ *respectively, let* $r_j$ *be non-negative integers and let* $a_j > 0$. *If* $\max\limits_{1\leq j\leq 2}\{m_j + r_j\} - \min\limits_{1\leq j\leq 2}\{l_j + r_j\} \leq 3$, *then the sequence* $\{a_1x_{i-r_1} + a_2y_{i-r_2}\}$ *is unimodal.*

**Example** 1. Suppose that $\{x_i\}_{i=0}^{3} = \{1, 3, 3, 2\}$ and $\{y_i\}_{i=1}^{4} = \{1, 2, 3, 3\}$. Obviously, they are unimodal and $l_1 = 1, m_1 = 2, l_2 = 3$ and $m_2 = 4$. Now consider the sequence $\{x_{i-1} + 2y_i\}$. Clearly, $r_1 = 1$ and $r_2 = 0$ and then $l = \min\{1 + 1, 3 + 0\} = 2$ and $m = \max\{2 + 1, 4 + 0\} = 4$. Since $m - l = 2 < 3$, $\{x_{i-1} + 2y_i\}$ is unimodal. In fact, $\{x_{i-1} + 2y_i\} = \{3, 7, 9, 8\}$ is clearly unimodal.

The following result is obtained by using the same technique in the argument of Theorem 2.1, which is its generalization.

**Theorem** 2.3 *Suppose that* $\{x_i^{(1)}\}_{i=q_1}^{n_1}$, $\{x_i^{(2)}\}_{i=q_2}^{n_2}$,$\cdots$,$\{x_i^{(k)}\}_{i=q_k}^{n_k}$ *are* $k$ *unimodal sequences of numbers for* $0 \leq q_j \leq n_j$, $1 \leq j \leq k$ *and* $k \geq 3$. *Let* $l_j, l_j+1, \cdots, m_j$ *be the modes of* $\{x_i^{(j)}\}$ *for* $1 \leq j \leq k$ *and* $l_j \leq m_j$, *let* $r_j$ *be non-negative integers and let* $a_j > 0$. *Then the sequence* $\{\sum\limits_{j=1}^{k} a_j x_{i-r_j}^{(j)}\}$ *is unimodal if and only if it is unimodal for* $\min\limits_{1\leq j\leq k}\{l_j + r_j\} \leq i \leq \max\limits_{1\leq j\leq k}\{m_j + r_j\}$.

If $m - l \leq 3$, then it is obvious that $\sum\limits_{j=1}^{k} a_j x_{n-r_j}^{(j)}$ is unimodal for $l \leq n \leq m$ and so the following conclusion holds.

**Corollary** 2.4 *Suppose that* $\{x_i^{(1)}\}_{i=q_1}^{n_1}$, $\{x_i^{(2)}\}_{i=q_2}^{n_2}$,$\cdots$,$\{x_i^{(k)}\}_{i=q_k}^{n_k}$ *are* $k$ *unimodal sequences of numbers for for* $0 \leq q_j \leq n_j$, $1 \leq j \leq k$ *and* $k \geq 3$. *Let* $l_j, l_j+1, \cdots, m_j$ *be the modes of* $\{x_i^{(j)}\}_{i=q_j}^{n_j}$ *for* $1 \leq j \leq k$ *and* $l_j \leq m_j$, *let* $r_j$ *be non-negative integers and let* $a_j > 0$. *If* $\max\limits_{1\leq j\leq k}\{m_j + r_j\} - \min\limits_{1\leq j\leq k}\{l_j + r_j\} \leq 3$, *then the sequence* $\{\sum\limits_{j=1}^{k} a_j x_{i-r_j}^{(j)}\}$ *is unimodal.*

Put $r_j = 0$ in Theorems 2.1, 2.3, Corollaries 2.2 and 2.4 and then one obtains the results as follows.

**Corollary** 2.5 *Suppose that* $\{x_i^{(1)}\}_{i=q_1}^{n_1}$, $\{x_i^{(2)}\}_{i=q_2}^{n_2}$,$\cdots$,$\{x_i^{(k)}\}_{i=q_k}^{n_k}$ *are* $k$ *unimodal sequences of numbers for for* $0 \leq q_j \leq n_j$, $1 \leq j \leq k$ *and* $k \geq 2$. *Let* $l_j, l_j+1, \cdots, m_j$ *be the modes of* $\{x_i^{(j)}\}$ *for* $1 \leq j \leq k$ *and* $l_j \leq m_j$ *and let* $a_j > 0$. *Then the sequence* $\{\sum\limits_{j=1}^{k} a_j x_i^{(j)}\}$ *is unimodal if and only if it is unimodal for* $\min\limits_{1\leq j\leq k} l_j \leq i \leq \max\limits_{1\leq j\leq k} m_j$.

**Corollary** 2.6 *Suppose that* $\{x_i^{(1)}\}_{i=q_1}^{n_1}$, $\{x_i^{(2)}\}_{i=q_2}^{n_2}$,$\cdots$,$\{x_i^{(k)}\}_{i=q_k}^{n_k}$ *are* $k$ *unimodal sequences of numbers for* $0 \leq q_j \leq n_j$, $1 \leq j \leq k$ *and* $k \geq 2$. *Let* $l_j, l_j+1, \cdots, m_j$ *be the modes of* $\{x_i^{(j)}\}$ *for*

$1 \leq j \leq k$ *and* $l_j \leq m_j$ *and let* $a_j > 0$. *If* $\max_{1\leq j\leq k} m_j - \min_{1\leq j\leq k} l_j \leq 3$, *then the sequence* $\{\sum_{j=1}^{k} a_j x_i^{(j)}\}$ *is unimodal.*

## 3. Unimodality of genus distributions for sets of ladder surfaces

Suppose that $a_l$ are distinct letters for $l \geq 1$. Let $R_1^n = a_{k_1}a_{k_2}a_{k_3}\cdots a_{k_r}$, $R_2^n = a_{k_{r+1}}a_{k_{r+2}}a_{k_{r+3}}\cdots a_{k_n}$, $R_3^n = a_{t_1}^- a_{t_2}^- a_{t_3}^- \cdots a_{t_s}^-$ and $R_4^n = a_{t_{s+1}}^- a_{t_{s+2}}^- a_{t_{s+3}}^- \cdots a_{t_n}^-$ where $n \geq k_1 > k_2 > k_3 > \cdots > k_r \geq 1$, $1 \leq k_{r+1} < k_{r+2} < k_{r+3} < \cdots < k_n \leq n$, $n \geq t_1 > t_2 > t_3 > \cdots > t_s \geq 1$, $1 \leq t_{s+1} < t_{s+2} < t_{s+3} < \cdots < t_n \leq n$ and $0 \leq r, s \leq n$, $k_p \neq k_q$, $t_p \neq t_q$ for $p \neq q$. The sets of ladder surfaces $\mathcal{S}_j^n$ are given below for $1 \leq j \leq 11$:

$$\mathcal{S}_1^n = \{R_1^n R_2^n R_3^n R_4^n\} \quad \mathcal{S}_2^n = \{R_1^n R_2^n R_4^n R_3^n\} \quad \mathcal{S}_3^n = \{R_1^n R_3^n R_2^n R_4^n\}$$

$$\mathcal{S}_4^n = \{aR_1^n R_2^n a^- R_3^n R_4^n\} \qquad \mathcal{S}_5^n = \{aR_1^n R_3^n a^- R_2^n R_4^n\}$$

$$\mathcal{S}_6^n = \{aR_1^n R_4^n a^- R_2^n R_3^n\} \qquad \mathcal{S}_7^n = \{aR_1^n a^- R_3^n R_2^n R_4^n\}$$

$$\mathcal{S}_8^n = \{R_1^n R_2^n a R_3^n a^- b R_4^n b^-\} \qquad \mathcal{S}_9^n = \{R_1^n R_3^n a R_2^n a^- b R_4^n b^-\}$$

$$\mathcal{S}_{10}^n = \{R_1^n R_4^n a R_2^n a^- b R_3^n b^-\} \qquad \mathcal{S}_{11}^n = \{R_1^n a R_2^n a^- b R_3^n b^- c R_4^n c^-\}$$

Let $g_{i_j}(n)$ denote the number of surfaces of genus $i$ in $\mathcal{S}_j^n$ for $1 \leq j \leq 11$ and $n \geq 1$. Suppose that $\gamma_j$ and $\gamma_{M_j}$ denote the minimum genus and the maximum genus of surfaces in $\mathcal{S}_j^n$. Then the *genus distribution* of $\mathcal{S}_j^n$ is the sequence of numbers $\{g_{i_j}(n)\}_{i=\gamma_j}^{\gamma_{M_j}}$:

$$g_{\gamma_j}(n), g_{\gamma_j+1}(n), \cdots, g_{\gamma_{M_j}}(n).$$

Since genus distributions of $\mathcal{S}_j^n$ are obviously log-concave and their modes are easily to obtained by Theorem 2.5 of [14] for $1 \leq j \leq 11$ and $1 \leq n \leq 3$, we sometimes omit them.

### 3.1 Log-concavity of genus distributions of $\mathcal{S}_j^n$ for $j = 1, 4, 6$

**Lemma 3.1**(Theorem 2.5 of [14]) *Suppose that* $g_{i_j}(n)$ *denotes the number of surfaces of genus* $i$ *in* $\mathcal{S}_j^n$ *for* $j = 1, 6$. *Let* $C_n(i) = \begin{pmatrix} n-i \\ i \end{pmatrix}$. *Then,*

$$g_{i_1}(n) = 2^{n+i}\frac{2n-3i}{n-i}C_n(i) \text{ for } 0 \leq i \leq \left\lceil \frac{n}{2} \right\rceil \text{ and } n \geq 1$$

and

$$g_{i_6}(n) = 2^{n+i-1}\frac{2n-3i+2}{n-i+1}C_{n+1}(i), \text{ if } 0 \leq i \leq \left\lceil \frac{n+1}{2} \right\rceil \text{ and } n \geq 1.$$

**Theorem 3.2** *Let $p_j(n)$ denote peaks of genus distributions of $\mathcal{S}_j^n$ for $j = 1, 6$ and $i \geq 1$. Genus distributions of $\mathcal{S}_j^n$ are log-concave for $j = 1, 6$ and $n \geq 1$. Then*

$$p_1(n) = \left[\frac{n+1}{3}\right], \text{ if } n \geq 1 \text{ and } n \neq 2 \tag{1}$$

*and*

$$p_6(n) = \left[\frac{n+2}{3}\right], \text{ if } n \geq 2. \tag{2}$$

**Proof.** By Lemma 3.1, for $1 \leq i \leq \frac{n}{2}$,

$$\frac{g_{i_1}(n)}{g_{(i-1)_1}(n)} = 2 \cdot \frac{n+1-2i}{i} \cdot \frac{n+2-2i}{n-i} \cdot \frac{2n-3i}{2n+3-3i}.$$

Since each item is non-increasing, $\frac{g_{i_1}(n)}{g_{(i-1)_1}(n)}$ is non-increasing and therefore the genus distribution of $\mathcal{S}_1^n$ is log-concave.

Now we compute $p_1(n)$. Put $n = 3m+k$ for $m \geq 0$ and $k = 0, 1, 2$. We verify the case $k = 2$ and leave others to readers. If $k = 2$, then $\left[\frac{n+1}{3}\right] = m+1$. By Lemma 3.1

$$\begin{aligned} g_{m_1}(n) &= 2^{3m+2+m} \frac{2(3m+2)-3m}{3m+2-m} C_{3m+2}(m) \\ &= 2^{4m+2}(3m+4) \frac{(2m+1)2m(2m-1)\cdots(m+3)}{m!}, \end{aligned}$$

$$\begin{aligned} g_{(m+1)_1}(n) &= 2^{3m+2+m+1} \frac{2(3m+2)-3(m+1)}{3m+2-(m+1)} C_{3m+2}(m+1) \\ &= 2^{4m+3}(3m+1) \frac{2m(2m-1)(2m-2)\cdots(m+2)}{m!} \end{aligned}$$

and

$$\begin{aligned} g_{(m+2)_1}(n) &= 2^{3m+2+m+2} \frac{2(3m+2)-3(m+2)}{3m+2-(m+2)} C_{3m+2}(m+2) \\ &= 2^{4m+4}(3m-2) \frac{(2m-1)(2m-2)\cdots(m-1)}{(m+2)!}. \end{aligned}$$

It is clear that

$$\frac{g_{(m+1)_1}(n)}{g_{m_1}} = \frac{6m^2+14m+4}{6m^2+11m+4} > 1$$

and that

$$\frac{g_{(m+1)_1}(n)}{g_{(m+2)_1}} = \frac{3m^2+7m+2}{3m^2-5m+2} > 1.$$

Thus $\left[\frac{n+1}{3}\right]$ is the peak of $\mathcal{S}_1^n$.

For $n \geq 2$, it is easily verified that for $0 \leq i \leq \left[\frac{n+1}{2}\right]$

$$g_{i_6}(n) = \frac{1}{4} g_{i_1}(n+1).$$

Thus the result holds for $\mathcal{S}_6^n$. $\square$

**Lemma** 3.3(Lemma 2.3 of [14]) *Let $g_{i_j}(n)$ be the number of surfaces in $S_n^j$ with genus $i$ for $j = 1$ and 4. Then for $n \geq 1$ and $1 \leq i \leq \left[\frac{n+1}{2}\right]$,*

$$g_{i_4}(n) = 4 g_{(i-1)_1}(n-1).$$

The following result is implied by Theorem 3.2 and Lemma 3.3.

**Corollary** 3.4 *Let $p_4(n)$ denote the modes of genus distribution of $\mathcal{S}_4^n$ for $n \geq 1$. The genus distribution of $\mathcal{S}_4^n$ is log-concave for each $n$ and*

$$p_4(n) = \begin{cases} 1, 2, & \text{if } n = 3; \\ \left[\frac{n}{3}\right] + 1, & \text{otherwise.} \end{cases}$$

### 3.2 Unimodality of genus distributions of $\mathcal{S}_j^n$ for $j = 2, 3, 7, 8, 10$

**Lemma 3.5**(Theorem 2.5 of [14]) *Let $g_{i_3}(n)$ be the number of surfaces in $S_n^3$ with genus $i$. Let $A_n(i) = \frac{2n-3i-2}{n-2i-1}$, let $B_n(i) = \frac{n-i-1}{n-2i}$ and let $C_n(i) = \begin{pmatrix} n-2-i \\ i \end{pmatrix}$. Then,*

$$g_{i_3}(n) = \begin{cases} 2^n + 4n - 2, & \text{if } i = 0 \text{ and } n \geq 1; \\ C_{n+2}(i+1)\Big(2^{3i+1} A_{n+2}(i+1) \\ \qquad + (2^{n+i-1} - 2^{3i-2}) \dfrac{(i+1) A_{n+2}(i) B_{n+2}(i+1)}{n-2i-1}\Big), \\ \qquad \text{if } 1 \leq i \leq \left[\frac{n}{2}\right] - 1 \text{ and } n \geq 2; \\ C_{n+1}(i)\Big(2^{3i+1} + (2^{n+i-1} - 2^{3i-2}) A_{n+2}(i) B_{n+2}(i+1)\Big), \\ \qquad \text{if } \left[\frac{n}{2}\right] - 1 < i \leq \left[\frac{n-1}{2}\right] \text{ and } n \geq 2; \\ (2^{n+i-1} - 2^{3i-2}) A_{n+2}(i) C_{n+2}(i), \\ \qquad \text{if } \left[\frac{n-1}{2}\right] < i \leq \left[\frac{n}{2}\right] \text{ and } n \geq 2. \end{cases}$$

**Lemma** 3.6(Lemma 2.3 of [14]) *Let $g_{i_j}(n)$ be the number of surfaces in $S_n^j$ with genus $i$ for $j = 3, 7, 10$ and $n \geq 0$. Let $f_{S_j^0}(x) = 1$. Then, for $n \geq 1$,*

$$g_{i_j}(n) = \begin{cases} g_{i_3}(n-1) + g_{i_6}(n-1) + 2g_{i_7}(n-1), \\ \qquad\qquad \text{if } j = 3, 0 \leq i \leq \left[\frac{n}{2}\right] \text{ and } n \geq 1; \\ 2g_{(i-1)_3}(n-1) + 2g_{i_{10}}(n-1), \\ \qquad\qquad \text{if } j = 7, 0 \leq i \leq \left[\frac{n+1}{2}\right] \text{ and } n \geq 1; \\ g_{(i-1)_6}(n-1) + 2g_{(i-1)_7}(n-1) + g_{i_{10}}(n-1), \\ \qquad\qquad \text{if } j = 10, 0 \leq i \leq \left[\frac{n}{2}\right] + 1 \text{ and } n \geq 1; \\ 0, \text{ otherwise.} \end{cases}$$

Lemmas $3.5 - 6$ imply the following result.

**Lemma** 3.7 *Let $g_{i_j}(n)$ be the number of surfaces in $S_j^n$ with genus $i$ for $j = 3, 7, 10$ and $n \geq 1$. Then*

$$g_{i_7}(n) = \begin{cases} 2, \text{ if } i = 0; \\ 4g_{(i-1)_3}(n-1) - 2, \text{ if } i = 1; \\ 4g_{(i-1)_3}(n-1), \text{ otherwise} \end{cases}$$

*and*

$$g_{i_{10}}(n) = \begin{cases} 1, \text{ if } i = 0; \\ g_{(i-1)_3}(n) - 1, \text{ if } i = 1; \\ g_{(i-1)_3}(n), \text{ otherwise.} \end{cases}$$

Therefore it is enough to find the unimodality of the genus distribution of $\mathcal{S}_3^n$ in order to study that of genus distributions of $\mathcal{S}_7^n$ and $\mathcal{S}_{10}^n$.

**Lemma** 3.8 *Let $g_{i_3}(n)$ be the number of surfaces in $S_3^n$ with genus $i$ and $n \geq 0$. Then for $n \geq 8$,*

$$g_{[\frac{n+2}{3}]_3}(n) > g_{([\frac{n+2}{3}]-1)_3}(n) \text{ and } g_{[\frac{n+2}{3}]_3}(n) > g_{([\frac{n+2}{3}]+1)_3}(n).$$

**Proof.** Put $n = 3m + k$ for $k = 0, 1, 2$. We verify the case $k = 0$ and leave others to readers. If $m = 3$, then by Lemma 3.5

$$g_{2_3}(9) = C_{9+2}(2+1)\Big(2^{6+1}A_{9+2}(2+1) + (2^{9+2-1} - 2^{6-2})\frac{(2+1)A_{9+2}(2)B_{9+2}(2+1)}{9-4-1}\Big) = 56432,$$

$$g_{3_3}(9) = C_{9+2}(3+1)\Big(2^{9+1}A_{9+2}(3+1) + (2^{9+3-1} - 2^{9-2})\frac{(3+1)A_{9+2}(3)B_{9+2}(3+1)}{9-6-1}\Big) = 126080$$

and

$$g_{4_3}(9) = C_{9+1}(4)\Big(2^{13} + (2^{9+4-1} - 2^{12-2})A_{9+2}(4)B_{9+2}(4+1)\Big) = 69632.$$

Thus

$$g_{3_3}(9) > g_{2_3}(9) \text{ and } g_{3_3}(9) > g_{4_3}(9).$$

For $m \geq 4$, by Lemma 3.5

$$\begin{aligned} g_{m_3}(3m) &= C_{3m+2}(m+1)\Big(2^{3m+1}A_{3m+2}(m+1) \\ &+ (2^{3m+m-1} - 2^{3m-2})\frac{(m+1)A_{3m+2}(m)B_{3m+2}(m+1)}{3m-2m-1}\Big) \\ &= \frac{(2m-1)(2m-2)\cdots(m+2)}{(m-1)!}\Big(2^{3m+1}(3m-1) \\ &+ (2^{4m} - 2^{3m-1})(3m+2)\Big) \end{aligned}$$

$$\begin{aligned} g_{(m-1)_3}(3m) &= C_{3m+2}(m-1+1)\Big(2^{3(m-1)+1}A_{3m+2}(m-1+1) \\ &+ (2^{3m+m-1-1} - 2^{3(m-1)-2})\frac{(m-1+1)A_{3m+2}(m-1)B_{3m+2}(m-1+1)}{3m-2(m-1)-1}\Big) \\ &= \frac{2m(2m-1)\cdots(m+2)}{(m-1)!}\Big(2^{3m-2}\frac{3m+2}{m+1} \\ &+ (2^{4m-2} - 2^{3m-5})\frac{m(3m+5)(2m+1)}{(m+3)(m+2)(m+1)}\Big) \end{aligned}$$

$$\begin{aligned} g_{(m+1)_3}(3m) &= C_{3m+2}(m+1+1)\Big(2^{3(m+1)+1}A_{3m+2}(m+1+1) \\ &+ (2^{3m+m+1-1} - 2^{3(m+1)-2})\frac{(m+1+1)A_{3m+2}(m+1)B_{3m+2}(m+1+1)}{3m-2(m+1)-1}\Big) \\ &= \frac{(2m-2)(2m-3)\cdots(m+3)}{(m-3)!}\Big(2^{3m+4}(3m-4) \\ &+ (2^{4m} - 2^{3m+1})\frac{(m+2)(3m-1)(2m-1)}{(m-3)(m-2)(m-1)}\Big) \end{aligned}$$

Then

$$\begin{aligned} \frac{g_{m_3}(3m)}{g_{(m-1)_3}(3m)} &= \frac{2^{3m+1}\frac{3m-1}{2m} + (2^{4m} - 2^{3m-1})\frac{(3m+2)}{2m}}{2^{3m-2}\frac{3m+2}{m+1} + (2^{4m-2} - 2^{3m-5})\frac{m(3m+5)(2m+1)}{(m+3)(m+2)(m+1)}} \\ &\geq \frac{2^{3m+1} + (3\cdot 2^{4m-1} - 3\cdot 2^{3m-2})}{3\cdot 2^{3m-2} + (3\cdot 2^{4m-1} - 3\cdot 2^{3m-4})} \\ &> 1 \end{aligned}$$

and

$$\begin{aligned}\frac{g_{m_3}(3m)}{g_{(m+1)_3}(3m)} &= \frac{(2m-1)(m+2)}{(m-2)(m-3)}\frac{2^{3m+1}(3m-1)+(2^{4m}-2^{3m-1})(3m+2)}{2^{3m+4}(3m-4)+(2^{4m}-2^{3m+1})\frac{(m+2)(3m-1)(2m-1)}{(m-3)(m-2)(m-1)}} \\ &> \frac{2^{3m+2}(3m-1)+(2^{4m+1}-2^{3m})(3m+2)}{2^{3m+4}(3m-4)+(3\cdot 2^{4m+1}-3\cdot 2^{3m+2})} \\ &> 1.\end{aligned}$$

Thus the result holds. □

**Theorem** 3.9 Let $p_3(n)$ denote the peak of the genus distribution of $\mathcal{S}_3^n$ for $n \geq 1$. Genus distributions of $\mathcal{S}_3^n$ are unimodal and

$$p_3(n) = \begin{cases} \left\lfloor \frac{n-1}{2} \right\rfloor, \text{ if } 1 \leq n \leq 6; \\ 2, \text{ if } n = 7; \\ \left\lceil \frac{n+2}{3} \right\rceil, \text{ if } n \geq 8. \end{cases}$$

**Proof.** The result is clear by Lemma 3.5 for $n \geq 8$. Next we verify the result by induction on $n(n \geq 8)$. Assume that the result holds for less than $n(n \geq 9)$.

Now we consider unimodality of the genus distribution of $\mathcal{S}_3^n$. By Lemma 3.6,

$$g_{i_3}(n) = g_{i_3}(n-1) + g_{i_6}(n-1) + 2g_{i_7}(n-1).$$

Here, $\mathcal{S}_3^{n-1}$ is unimodal and $p_3(n-1) = \left\lceil \frac{n+1}{3} \right\rceil$ by the induction hypothesis. $\mathcal{S}_6^{n-1}$ is unimodal and $p_6(n-1) = \left\lceil \frac{n+1}{3} \right\rceil$ by Theorem 3.2. Clearly, $\mathcal{S}_7^{n-1}$ is unimodal and $p_7(n-1) = \left\lfloor \frac{n}{3} \right\rfloor + 1$ by Lemma 3.7 and induction hypothesis. Then by applying Corollary 2.6, $\mathcal{S}_3^n$ is unimodal. Combining with Lemma 3.8, we get

$$p_3(n) = \left\lceil \frac{n+2}{3} \right\rceil$$

as desired. □

Armed with Theorem 3.9 and Lemma 3.7, the following conclusion is easily induced.

**Corollary** 3.10 Let $p_j(n)$ denote the peak of genus distribution of $\mathcal{S}_j^n$ for $j = 7, 10$ and $n \geq 1$. Genus distributions of $\mathcal{S}_j^n$ are unimodal and

$$p_7(n) = \begin{cases} \left\lceil \frac{n}{2} \right\rceil, \text{ if } 2 \leq n \leq 7; \\ 3, \text{ if } n = 8; \\ \left\lceil \frac{n+1}{3} \right\rceil + 1, \text{ if } n \geq 9 \end{cases}$$

and

$$p_{10}(n) = \begin{cases} \left[\frac{n+1}{2}\right], \text{ if } 1 \leq n \leq 6; \\ 3, \text{ if } n = 7; \\ \left[\frac{n+2}{3}\right] + 1, \text{ if } n \geq 8. \end{cases}$$

By Lemma 2.3 of [14] for $n \geq 1$

$$g_{i_j}(n) = \begin{cases} 4g_{i_7}(n-1), \text{ if } j = 2, 0 \leq i \leq \left[\frac{n}{2}\right]; \\ 4g_{(i-1)_7}(n-1), \text{ if } j = 8, 1 \leq i \leq \left[\frac{n}{2}\right] + 1. \end{cases}$$

Thus

**Corollary** 3.11 Let $p_j(n)$ denote the peak of the genus distribution of $\mathcal{S}_j^n$ for $j = 2, 8$ and $n \geq 3$. Then genus distributions of $\mathcal{S}_j^n$ are unimodal and

$$p_2(n) = \begin{cases} \left[\frac{n-1}{2}\right], \text{ if } 3 \leq n \leq 8; \\ 3, \text{ if } n = 9; \\ \left[\frac{n}{3}\right] + 1, \text{ if } n \geq 10 \end{cases}$$

and

$$p_8(n) = \begin{cases} \left[\frac{n+1}{2}\right], \text{ if } 3 \leq n \leq 8; \\ 4, \text{ if } n = 9; \\ \left[\frac{n}{3}\right] + 2, \text{ if } n \geq 10. \end{cases}$$

### 3.3 Unimodality of genus distributions of $\mathcal{S}_j^n$ for $j = 5, 9, 11$

**Lemma 3.12**(Theorem 2.5 of [14]) *Let* $g_{0_5}(1) = 2$, $g_{1_5}(1) = 2$, $g_{0_5}(2) = 2$, $g_{1_5}(2) = 14$, $g_{0_9}(1) = 1$, $g_{1_9}(1) = 3$, $g_{1_9}(2) = 10$, $g_{2_9}(2) = 6$, $g_{1_9}(3) = 10$, $g_{2_9}(3) = 54$, $B_n(i) = \frac{n-i-1}{n-2i}$, $C_n(i) = \begin{pmatrix} n-2-i \\ i \end{pmatrix}$ *and* $D_n(i) = \frac{n}{i}2^i$. *Then,* $g_{i_j}(n) =$

$$\begin{cases} 2^n + 8n + 8, \text{ if } j = 5, i = 1 \text{ and } n = 3, 4; \\ 2^n + 8n, \text{ if } j = 5, i = 1 \text{ and } n \geq 5; \\ (2^n - 2^{2i-2})C_n(i-2)D_n(i-1) + 2^{2i}C_n(i-1)D_n(i), \\ \qquad \text{if } j = 5, 2 \leq i < \frac{n}{2} - 1 \text{ and } n \geq 5; \\ (2^n - 2^{2i-2})C_n(i-2)D_n(i-1) + 2^{2i}C_n(i-1)D_n(i) + 2^{n-1}, \\ \qquad \text{if } j = 5, i = \frac{n}{2} - 1 \text{ and } n \geq 5; \end{cases}$$

$$
\begin{aligned}
&(2^n - 2^{2i-2})C_n(i-2)D_n(i-1) + 2^{2i}C_n(i-1)D_n(i) + 2^n,\\
&\qquad\qquad \textit{if } j = 5, \frac{n}{2} - 1 < i \le \frac{n-1}{2} \textit{ and } n \ge 4;\\
&(2^n - 2^{2i-2})C_n(i-2)D_n(i-1) + 2^{\frac{3n}{2}+1} - 3\cdot 2^{n-1},\\
&\qquad\qquad \textit{if } j = 5, \frac{n-1}{2} < i \le \frac{n}{2} \textit{ and } n \ge 4;\\
&(2^n - 2^{2i-2})C_n(i-2)D_n(i-1),\\
&\qquad\qquad \textit{if } j = 5, \frac{n}{2} < i \le \frac{n+1}{2} \textit{ and } n \ge 3;\\
&6, \textit{ if } j = 9, i = 1 \textit{ and } n \ge 4;\\
&3\cdot 2^n + 48n - 86, \textit{ if } j = 9, i = 2 \textit{ and } n = 4, 5;\\
&3\cdot 2^n + 48n - 102, \textit{ if } j = 9, i = 2 \textit{ and } n \ge 6;\\
&3C_n(i-1)\Big(2^{3i-2}B_{n+1}(i)\\
&\qquad +(2^{n+i-2} - 2^{3i-5})\frac{(i-1)B_n(i-1)B_{n+1}(i-1)}{n-2i+1}\Big),\\
&\qquad\qquad \textit{if } j = 9, 3 \le i < \frac{n-1}{2} \textit{ and } n \ge 6;\\
&3C_n(i-1)\Big(2^{3i-2}B_{n+1}(i)\\
&\qquad +(2^{n+i-2} - 2^{3i-5})\frac{(i-1)B_n(i-1)B_{n+1}(i-1)}{n-2i+1}\Big) + 2^{n-1},\\
&\qquad\qquad \textit{if } j = 9, i = \frac{n-1}{2} \textit{ and } n \ge 7;\\
&3C_n(i-1)\Big(2^{3i-2}B_{n+1}(i)\\
&\qquad +(2^{n+i-2} - 2^{3i-5})\frac{(i-1)B_n(i-1)B_{n+1}(i-1)}{n-2i+1}\Big) + 2^n,\\
&\qquad\qquad \textit{if } j = 9, \frac{n-1}{2} < i \le \frac{n}{2} \textit{ and } n \ge 6;\\
&C_{n-1}(i-2)\Big(2^{3i-2} + 3(2^{n+i-2} - 2^{3i-5})B_n(i-1)B_{n+1}(i-1)\Big)\\
&\qquad +2^{\frac{3n+1}{2}} - 3\cdot 2^{n-1},\\
&\qquad\qquad \textit{if } j = 9, \frac{n}{2} < i \le \frac{n+1}{2} \textit{ and } n \ge 5;\\
&3(2^{n+i-2} - 2^{3i-5})B_{n+1}(i-1)C_n(i-2),\\
&\qquad\qquad \textit{if } j = 9, \frac{n+1}{2} < i \le \frac{n}{2} + 1 \textit{ and } n \ge 4.
\end{aligned}
$$

Applying a similar way in the argument of Lemma 3.8, the following conclusion holds.

**Lemma** 3.13 *Let $g_{i_j}(n)$ be the number of surfaces in $S_j^n$ of genus $i$ for $j = 5, 9$ and $n \ge 6$. Then*

$$g_{([\frac{n+1}{3}]+1)_5}(n) > g_{[\frac{n+1}{3}]_5}(n) \textit{ and } g_{([\frac{n+1}{3}]+1)_5}(n) > g_{([\frac{n+1}{3}]+2)_5}(n) \textit{ for } 6 \le n \le 16,$$

$$g_{([\frac{n+2}{3}]+1)_5}(n) > g_{[\frac{n+2}{3}]_5}(n) \textit{ and } g_{([\frac{n+2}{3}]+1)_5}(n) > g_{([\frac{n+2}{3}]+2)_5}(n) \textit{ for } n \ge 17$$

*and*

$$g_{([\frac{n}{3}]+2)_9}(n) > g_{([\frac{n}{3}]+1)_9}(n) \text{ and } g_{([\frac{n}{3}]+2)_9}(n) > g_{([\frac{n}{3}]+3)_9}(n) \text{ for } n \geq 10.$$

**Lemma** 3.14(Lemma 2.3 of [14]) *Let $g_{i_j}(n)$ be the number of surfaces in $S_j^n$ with genus $i$ for $j = 5, 9, 11$ and $n \geq 0$. Let $f_{S_j^0}(x) = 1$. Then, for $n \geq 1$,*

$$g_{i_j}(n) = \begin{cases} 2g_{(i-1)_3}(n-1) + 2g_{i_9}(n-1), \\ \qquad \text{if } j = 5, 0 \leq i \leq \left[\frac{n+1}{2}\right] \text{ and } n \geq 1; \\ g_{(i-1)_5}(n-1) + 2g_{(i-1)_7}(n-1) + g_{i_{11}}(n-1), \\ \qquad \text{if } j = 9, 1 \leq i \leq \left[\frac{n}{2}\right] + 1 \text{ and } n \geq 1; \\ 2g_{(i-1)_9}(n-1) + 2g_{(i-1)_{10}}(n-1), \\ \qquad \text{if } j = 11, 1 \leq i \leq \left[\frac{n+1}{2}\right] + 1 \text{ and } n \geq 1; \\ 0, \text{ otherwise.} \end{cases}$$

**Theorem 3.15** Let $p_9(n)$ denote the peak of genus distribution of $\mathcal{S}_9^n$ for $n \geq 2$. Genus distributions of $\mathcal{S}_9^n$ are unimodal and

$$p_9(n) = \begin{cases} \left[\frac{n+1}{2}\right], \text{ if } 2 \leq n \leq 8; \\ 4, \text{ if } n = 9; \\ \left[\frac{n}{3}\right] + 2, \text{ if } n \geq 10 \end{cases}$$

**Proof.** We verify the conclusion by induction on $n$. It is obvious for $n = 2$. Assume that it holds for less than $n (n \geq 3)$.

By Lemma 3.14, for $1 \leq i \leq \left[\frac{n}{2}\right] + 1$ and $n \geq 3$,

$$g_{i_9}(n) = g_{(i-1)_5}(n-1) + 2g_{(i-1)_7}(n-1) + 2g_{(i-1)_9}(n-2) + 2g_{(i-1)_{10}}(n-2). \tag{3}$$

Here, genus distributions of $\mathcal{S}_7^{n-1}$ and $\mathcal{S}_{10}^{n-2}$ are unimodal by Corollary 3.10. The genus distritution of $\mathcal{S}_9^{n-1}$ is unimodal by the induction hypothesis. Now consider unimodality of the genus distribution of $\mathcal{S}_5^{n-1}$. By Lemma 3.14, for $1 \leq i \leq \left[\frac{k}{2}\right]$ and $k \geq 2$,

$$g_{i_5}(k-1) = 2g_{(i-1)_3}(k-2) + 2g_{i_9}(k-2)$$

where the genus distribution of $\mathcal{S}_3^{n-2}$ is unimodal by Theorem 3.9 and where that of $\mathcal{S}_9^{n-2}$ is unimodal according to the induction hypothesis. Since it is clear that $p_9(n-2) - (p_3(n-2)+1) \leq$

3, that of $\mathcal{S}_5^{n-1}$ is unimodal by Corollary 2.2. Combining with Lemma 3.13, we get for $k \leq n$

$$p_5(k-1) = \begin{cases} \left[\frac{k-1}{2}\right], \text{ if } 3 \leq k \leq 6; \\ \left[\frac{k}{3}\right] + 1, \text{ if } 7 \leq k \leq 17; \\ \left[\frac{k+1}{3}\right] + 1, \text{ if } k \geq 18. \end{cases}$$

Then we consider the unimodality of genus distribution of $\mathcal{S}_9^n$. It is easily known that

$$p_7(n-1) \leq p_9(n-2) \leq p_{10}(n-2) \leq p_5(n-1) \text{ and } p_5(n-1) + 1 - (p_7(n-1)+1) \leq 3. \quad (4)$$

Thus the conclusion is true for $n$ by armed with $(3-4)$, Corollary 2.4 and Lemma 3.13 and so is the conclusion by induction. $\square$

By a similar way in the argument of Theorem 3.9, the following result is obtained.

**Theorem** 3.16 Let $p_5(n)$ denote the peak of genus distribution of $\mathcal{S}_5^n$ for each $n \geq 2$. The genus distribution of $\mathcal{S}_5^n$ is unimodal and

$$p_5(n) = \begin{cases} \left[\frac{n}{2}\right], \text{ if } 2 \leq n \leq 5; \\ \left[\frac{n+1}{3}\right] + 1, \text{ if } 6 \leq n \leq 16; \\ \left[\frac{n+2}{3}\right] + 1, \text{ if } n \geq 17. \end{cases}$$

Since Lemma 3.12 implies for $n \geq 3$

$$g_{i_{11}}(n) = \begin{cases} 2, \text{ if } i = 1; \\ g_{(i-1)_5}(n) - 2, \text{ if } i = 2; \\ g_{(i-1)_5}(n), \text{ otherwise}, \end{cases}$$

it is easy to get the following result.

**Corollary** 3.17 Let $p_{11}(n)$ denote the peak of the genus distribution of $\mathcal{S}_j^n$ for each $n \geq 2$. Then the genus distribution of $\mathcal{S}_{11}^n$ is unimodal and

$$p_{11}(n) = \begin{cases} \left[\frac{n}{2}\right] + 1, \text{ if } 2 \leq n \leq 5; \\ \left[\frac{n+1}{3}\right] + 2, \text{ if } 6 \leq n \leq 16; \\ \left[\frac{n+2}{3}\right] + 2, \text{ if } n \geq 17. \end{cases}$$

## 4. Unimodality of genus distribution of ladders and crosses

Let $e_0$ and $e_1$ be edges of a connected graph $G_0$. Add vertices $u_1, u_2, \cdots, u_n$ on $e_0$ and add vertices $v_1, v_2, \cdots, v_n$ in sequence for $n \geq 1$. If one adds $u_l v_l$ denoted by $a_l$ such that they are parallel for $1 \leq l \leq n$, then a ladder $GL_n$ is constructed. Otherwise one adds $u_l v_{n-l+1}$ and then a cross $GC_n$ is obtained(See Fig.1(a) and (b)).




$GL_n$ (a) $GC_n$ (b)

Fig.1: $GL_n$ and $GC_n$

**Lemma** 4.1 (Theorem 3.1 of [14]) *Let $f_G(x)$ denote the genus polynomial of a graph $G$. Then*

$$f_{GL_n}(x) = \sum_{j=1}^{11} f_j(x) f_{\mathcal{S}_j^n}(x)$$

*where* $f_G(x) = \sum\limits_{i=1}^{11} f_j(x)$.

Similarly, the result is clear for a cross $GC_n$.

**Lemma** 4.2 *Let $f_G(x)$ denote the genus polynomial of a graph $G$. Then*

$$f_{GC_n}(x) = \sum_{j=1}^{11} f_j(x) f_{\mathcal{S}_j^n}(x)$$

*where* $f_G(x) = \sum\limits_{i=1}^{11} f_j(x)$.

Based on Lemmas $4.1 - 2$, unimodality of genus distributions of some ladders and crosses can be determined by using criteria in Section 2.

**Lemma** 4.3 (Propositions $2 - 3$ of [10]) *Let $a(x)$ and $b(x)$ be polynomials with positive coefficients.*

(1) *If both $a(x)$ and $b(x)$ are log-concave, then so is $a(x)b(x)$.*

(2) *If both $a(x)$ is log-concave and $b(x)$ is unimodal, then $a(x)b(x)$ is unimodal.*

Armed with Lemma 4.3, unimodality of genus distributions of some ladders and crosses is determined as follows.

**Theorem** 4.4 *Let $G_n$ be a ladder or a cross. Suppose that $f_{\mathcal{P}_{G_n}}(x) = f_{\mathcal{P}_{G_0}}(x) f_{\mathcal{S}_j^n}(x)$ for $j = 1$ or 6. If $f_{\mathcal{P}_{G_0}}(x)$ is log-concave(or unimodal), then $f_{\mathcal{P}_{G_n}}(x)$ is log-concave (or unimodal).*

Next we consider several types of ladders which are Closed-end ladders $L_n$, circular ladders $CL_n$, Möbius ladders $ML_n$, Ringel ladders $RL_n$ and a type of crosses $R_n$. See Fig.2 for $n = 4$.




Fig.2:$L_4$,$CL_4$,$ML_4$ and $R_4$

By applying Section 3 of [14],

$$f_{L_n}(x) = f_{S_6^n}(x).$$

Then the following conclusion is immediate by Theorem 3.2, which induces the known results for Closed-end ladders in [3].

**Corollary** 4.5 *Let $p_L(n)$ denote the peak of the genus distribution of a Closed-end ladder $L_n$ for each $n \geq 1$. The genus distribution of $L_n$ is log-concave and*

$$p_L(n) = \left\lceil \frac{n+2}{3} \right\rceil, \text{ if } n \geq 2.$$

For $CL_n$ and $ML_n$, since genus distribution for $ML_n$ equals to that of $CL_n$, except that $ML_n$ has two extra embeddings of genus 1 and two fewer embeddings of genus 0([8]), it is enough to consider unimodality of the genus distribution for $CL_n$. In [14] we have

$$g_i(CL_n) = 2g_{i_9}(n-1) + 2g_{i_{10}}(n-1).$$

Since by Lemma 3.14

$$g_{i_{11}}(n) = 2g_{(i-1)_9}(n-1) + 2g_{(i-1)_{10}}(n-1),$$

$$g_i(CL_n) = g_{(i+1)_{11}}(n).$$

Thus, we have the following result by Corollary 3.17.

**Corollary** 4.6 *Let $p_{CL}(n)$ and $p_{ML}(n)$ denote the peaks of genus distributions of circular ladders $CL_n$ and Möbius ladders $ML_n$ for $n \geq 2$,respectively. Then their genus distributions are unimodal and*

$$p_{CL}(n) = p_{ML}(n) = \begin{cases} \left\lceil \frac{n}{2} \right\rceil, \text{ if } 2 \leq n \leq 5; \\ \left\lceil \frac{n+1}{3} \right\rceil + 1, \text{ if } 6 \leq n \leq 16; \\ \left\lceil \frac{n+2}{3} \right\rceil + 1, \text{ if } n \geq 17. \end{cases}$$

Similarly, since in [14]

$$g_i(RL_n) = 2g_{(i-1)_3}(n) + 2g_{i_{10}}(n)$$

and since by Lemma 3.6

$$g_{i_7}(n+1) = 2g_{(i-1)_3}(n) + 2g_{i_{10}}(n),$$

the following result is implied by Corollary 3.10.

**Corollary** 4.7 *Let $p_{RL}(n)$ denote the peak of the genus distribution of a Ringel ladder $RL_n$ for each $n \geq 1$. Then its genus distribution is unimodal and*

$$p_{RL}(n) = \begin{cases} \left\lceil \frac{n+1}{2} \right\rceil, \text{ if } 1 \leq n \leq 6; \\ 3, \text{ if } n = 7; \\ \left\lceil \frac{n+1}{3} \right\rceil + 1, \text{ if } n \geq 8 \end{cases}$$

Now consider the type of crosses $R_n$. The following equation is immediate from [13] and Theorem 3 in [16].

$$g_i(R_n) = 2\mu_{i_6}(n) + 2\mu_{(i-1)_1}(n) = 2g_{i_5}(n) + 2g_{(i-1)_2}(n).$$

By applying the same technique in the argument of Theorem 3.9, the following result is obtained.

**Theorem** 4.8 *Let $p_R(n)$ denote the peak of the genus distribution of $R_n$ for each $n \geq 1$. Then its genus distribution is unimodal and*

$$p_R(n) = \begin{cases} \left\lceil \frac{n+1}{2} \right\rceil, \text{ if } 1 \leq n \leq 6; \\ 3, \text{ if } n = 7; \\ \left\lceil \frac{n+2}{3} \right\rceil + 1, \text{ if } n \geq 8. \end{cases}$$

## 5. Further study

**Problem** 5.1 *Determine whether genus distributions of sets of ladder surfaces $\mathcal{S}_j^n$ are log-concave for $2 \leq j \leq 11$, $j \neq 4, 6$ and $n \geq 2$.*

**Problem** 5.2 *Let $k$ sequences of polynomials $\{P_j(n)\}$ satisfy certain dependent recurrence relations for $k \geq 2$ and $1 \leq j \leq k$. If their explicit expressions are unknown, then determine whether they are unimodal (or log-concave). For example, let $P_1(n) = \sum_{i=0}^{n} g_i(n)x^i$ and let $P_2(n) = \sum_{i=1}^{n} g_{i_2}(n)x^i$ where $g_0(0) = 1$, $g_0(1) = 2$, $g_1(1) = 14$, $g_{1_2}(1) = 4$ (Theorem 4.1 [15]). For $n \geq 2$,*

$$\begin{cases} g_i(n) = 2g_i(n-1) + 8g_{i-1}(n-1) + 48g_{i-1}(n-2) + 12g_{(i-1)_2}(n-1), \\ g_{i_2}(n) = 8g_{(i-1)_2}(n-1) + 32g_{i-1}(n-2). \end{cases}$$

determine whether $\{P_1(n)\}$ and $\{P_2(n)\}$ are unimodal ( or log-concave ).

**Problem** 5.3 *Suppose that $\{x_i\}_{i=q_1}^{n_1}$ and $\{y_i\}_{i=q_2}^{n_2}$ are unimodal sequences of numbers for $0 \leq q_1 \leq n_1$ and $0 \leq q_2 \leq n_2$. Let $l_1, l_1+1, \cdots, m_1$ and $l_2, l_2+1, \cdots, m_2$ be the modes of $\{x_i\}_{i=q_1}^{n_1}$ and $\{y_i\}_{i=q_2}^{n_2}$ for $j = 1, 2$ and $l_j \leq m_j$ respectively, let $r_j$ be non-negative integers and let $a_j > 0$. If $\max_{1\leq j\leq 2}\{m_j + r_j\} - \min_{1\leq j\leq 2}\{l_j + r_j\} \geq 4$, then determine the conditions such that the sequence of numbers $\{a_1x_{i-r_1} + a_2y_{i-r_2}\}$ is unimodal.*

**Problem** 5.4 *Suppose that $\{x_i^{(1)}\}_{i=q_1}^{n_1}$, $\{x_i^{(2)}\}_{i=q_2}^{n_2}, \cdots, \{x_i^{(k)}\}_{i=q_k}^{n_k}$ are $k$ unimodal sequences of numbers for $k \geq 3$. Let $l_j, l_j+1, \cdots, m_j$ be the modes of $\{x_i^{(j)}\}$ for $1 \leq j \leq k$ and $l_j \leq m_j$, let $r_j$ be non-negative integers and let $a_j > 0$. If $\max_{1\leq j\leq 2}\{m_j + r_j\} - \min_{1\leq j\leq 2}\{l_j + r_j\} \geq 4$, then determine the conditions such that the sequence of numbers $\{\sum_{j=1}^{k} a_j x_{i-r_j}^{(j)}\}$ is unimodal.*

# References


[1] F. Brenti, Log-concave and unimodal sequences in algebra, combinatorics, and geometry: An update, Contemp. Math., 178(1994) 71-89.

[2] Y. Chen, A note on a conjecture of Stahl, Canad. J. Math 60(2008) 958-959.

[3] M.L. Furst, J.L. Gross, R. Statman, Genus distributions for two classes of graphs, J. Combin. Theory(B) 46(1989) 22-36.

[4] J.L.Gross, T. Mansour and T. W. Tucker, Log-concavity of genus distributions of ring-like families of graphs, submitted.

[5] J.L. Gross, D.P. Robbins, T.W. Tucker, Genus distributions for bouquets of circles, J. Combin. Theory(B) 47(1989) 292-306.

[6] Y.P. Liu, *Advances in Combinatorial Maps* (in Chinese), Northern Jiaotong University Press, Beijing, 2003.

[7] L. Liu, Y. Wang, A unified approach to polynomial sequences with only real zeros, Advances in Applied Mathematics 38 (2007) 542-560.

[8] L. A. McGeoch, Algorithms for two graph problems: computing maximum-genus imbeddings and the two-server problem, Ph.D Thesis, Computer Science Dept., Carnegie Mellon University, PA, 1987.

[9] S. Stahl, On the zeros of some genus polynomials,Can. J. Math. 49(1997) 617-640.

[10] R.P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, Ann. New York Acad. Sci. 576 (1989) 500-535.

[11] E. H. Tesar, Genus distribution of Ringel ladders, Discrete Math. 216(2000) 235-252.

[12] C. Thomassen, The graph genus problem is NP-complete, J. Algorithms 10(1989) 568-576.

[13] L.X. Wan, Y.P. Liu, Orientable embedding distributions by genus for certain type of non-planar graphs (I), Ars Combin. 79(2006) 97-105.

[14] L.X. Wan, Genus Distritution of General Ladders (in chinese),Acta Math. Appl. Sin.,31(2008) 806-816.

[15] L.X. Wan, Y.P. Liu, *Orientable embedding distribution for certain types of graphs*, J. Combin. Theory(B) 98(2008) 19-32.

[16] L.X. Wan, Y.P. Liu, On embedding genus distribution of ladders and crosses, Applied Math. Letters 22(2009) 738-742.

[17] X.M. Zhao, Y.P. Liu, Unimodality of Genus Distribution for Tree-Graph, J. Shanxi Univ. (Nat. Sci. Ed. ) 29(2006) 242-244.